



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

V. *The Substance of some Experiments of planting Seeds in Moss, lately made by Mr. Charles Bonnet, of Geneva, F. R. S.*

Read Feb. 18. 1747-8. **M**R. Bonnet was inclined to try whether Plants were capable of Vegetation, when they were only set in Moss, instead of being planted in the Earth.

With this Design, he filled with Moss several Garden Pots, and he compressed the Moss more or less, as he judged, the several Plants he intended to place in them, might respectively require a closer or a looser Soil.

He then sowed in Moss, Wheat, Barley, Oats, and Pease. And he found, first, that all the Grains sowed in that manner came to Maturity later than those of the same Sorts which were sowed at the same time in Mould.

2dly. That the Stems from the several Grains sowed in the Moss were generally taller than those which sprung from the Ground.

3dly. There came from the Grains sowed in the Moss a greater Number of Blades than from the Grains sowed in the Earth.

4thly. The Grains sowed in Moss produced more plentifully than the others.

5thly. Those Grains that were gather'd, from the Produce of those which vegetated in the Moss, having been again sowed some in Moss. and some in Earth, succeeded well in both.

Mr. Bonnet has also planted in Moss, Pinks, Gilly-flowers, Daisies, Tuberoses, Tulips, Hyacinths, Jonquils, and Narcissus's; and all these Plants succeeded as well

well as others of the same Sorts, which he at the same time planted in Mould.

He also placed in Moss Cuttings and Layers of Vines, and these Cuttings and Layers became Vines; and these Vines in a short time grew larger than others, that came from Cuttings and Layers planted at the same time in the Ground.

VI. *A Continuation of an Account of an Essay towards a Natural History of Carolina, and the Bahama Islands; by Mark Catesby, F. R. S. with some Extracts out of the Appendix, * by C. Mortimer, Secret. R. S.*

Read Feb. 18. 1747-8. *U* *Regallus minor, fuscus; cervice plumis alas imitantibus donatâ.*

This Bird was about a third Part bigger than a common Partridge, all over of a reddish Brown, marked transversely with black and white waved Lines intermixed; but what is singular and extraordinary in this Bird, and distinguishes it from all others yet known, are two Tufts of Feathers resembling little Wings, 3 Inches long, placed on the hind Part of the Head, opposite to one another. These little Wings (if so they may be called) were fixed, as our Author says, to the Neck, in like manner as the real Wings are to the Body; whereby it had the like Power of contracting and dilating them; and they may

X 2

possibly